

Numoisyn Lozenges

Prescribing Information

Ingredients: Sorbitol (0.3 g per lozenge), polyethylene glycol, malic acid, sodium citrate, calcium phosphate dibasic, hydrogenated cottonseed oil, citric acid, magnesium stearate, and silicon dioxide.

Pharmaceutical Form: Oral lozenge

Contents: 100 lozenges per bottle. Net weight of 40 g (0.4 g per lozenge).

Therapeutic Group: Numoisyn Lozenges are oral lozenges formulated to promote lubrication of oral mucosa that may be dry due to a variety of circumstances, including medication, chemotherapy or radiotherapy, Sjögren's syndrome, or oral inflammation.

Indications: Numoisyn Lozenges are indicated for the treatment of xerostomia (dry mouth). Numoisyn Lozenges provide temporary relief of dry mouth due to damaged salivary function. Numoisyn Lozenges are formulated to support the natural protection of teeth provided by saliva so that no damage occurs to teeth with repeated use of the lozenges.

Contraindications: Numoisyn Lozenges are contraindicated in patients with fructose intolerance or a known history of hypersensitivity to any of the ingredients.

Warning: Federal law restricts Numoisyn Lozenges to sale by, or on the order of, a physician or properly licensed practitioner.

Interactions: There are no known interactions between Numoisyn Lozenges and any medicinal or other products.

Directions for Use: Let one Numoisyn Lozenge dissolve slowly in the mouth when needed. To obtain optimal effect, move the lozenge around in the mouth. Repeat as necessary. Do not exceed 16 lozenges in 24 hours.

Side Effects: Excessive consumption can cause minor digestive problems.

Storage: Store at room temperature. KEEP OUT OF REACH OF CHILDREN.

Overdose: No overdoses have been reported to date.

Manufactured in Italy under license from
Sinclair Pharmaceuticals Ltd.
Godalming, Surrey GU7 1XW UK

Distributed by
ALIGN Pharmaceuticals, LLC
Berkeley Heights, NJ 07922 USA

www.alignpharma.com

